

Research Article

# Cell-free protein synthesis enables high yielding synthesis of an active multicopper oxidase

Jian Li<sup>1,2,3,4</sup>, Thomas J. Lawton<sup>5</sup>, Jan S. Kostecki<sup>6</sup>, Alex Nisthal<sup>6</sup>, Jia Fang<sup>5</sup>, Stephen L. Mayo<sup>6</sup>, Amy C. Rosenzweig<sup>3,5,7</sup> and Michael C. Jewett<sup>1,2,3,4</sup>

<sup>1</sup> Department of Chemical and Biological Engineering, Northwestern University, Evanston, IL, USA

<sup>2</sup> Chemistry of Life Processes Institute, Northwestern University, Evanston, IL, USA

<sup>3</sup> Robert H. Lurie Comprehensive Cancer Center, Northwestern University, Chicago, IL, USA

<sup>4</sup> Simpson Querrey Institute for BioNanotechnology in Medicine, Northwestern University, Chicago, IL, USA

<sup>5</sup> Department of Molecular Biosciences, Northwestern University, Evanston, IL, USA

<sup>6</sup> Division of Biological Sciences, California Institute of Technology, Pasadena, CA, USA

<sup>7</sup> Department of Chemistry, Northwestern University, Evanston, IL, USA

Multicopper oxidases (MCOs) are broadly distributed in all kingdoms of life and perform a variety of important oxidative reactions. These enzymes have potential biotechnological applications; however, the applications are impeded by low expression yields in traditional recombinant hosts, solubility issues, and poor copper cofactor assembly. As an alternative to traditional recombinant protein expression, we show the ability to use cell-free protein synthesis (CFPS) to produce complex MCO proteins with high soluble titers. Specifically, we report the production of MCOs in an *Escherichia coli*-based cell-free transcription-translation system. Total yields as high as 1.2 mg mL<sup>-1</sup> were observed after a 20-h batch reaction. More than 95% of the protein was soluble and activity was obtained by simple post-CFPS addition of copper ions in the form of CuSO<sub>4</sub>. Scale-up reactions were achieved from 15 to 100 μL without a decrease in productivity and solubility. CFPS titers were higher than in vivo expression titers and more soluble, avoiding the formation of inclusion bodies. Our work extends the utility of the cell-free platform to the production of active proteins containing copper cofactors and demonstrates a simple method for producing MCOs.

Received	22 APR 2015
Revised	26 JUN 2015
Accepted	31 JUL 2015
Accepted article online	03 AUG 2015

**Keywords:** Cell-free protein synthesis · Metalloprotein · Multicopper oxidase · Transcription and translation

## 1 Introduction

Cell-free protein synthesis (CFPS) is emerging as a robust platform for fundamental and applied research in the areas of synthetic biology and biotechnology [1, 2]. CFPS has been developed with several different organisms (e.g. *E. coli*, wheat germ, rabbit reticulocytes, CHO cells, yeast, and insect cells) to produce various proteins including therapeutic vaccines, virus-like particles, and membrane

proteins [3–26]. CFPS is also being used for high-throughput functional and structural studies of a wide variety of proteins [27, 28], as well as more recently for rapid prototyping of biological circuits [29–35]. Although different CFPS systems have their own advantages, the *E. coli*-based system is the most commonly used due to fast cell growth of the source strain, inexpensive and streamlined extract preparation, and generally high protein yields. Because of the absence of cell walls in cell-free systems, cofactors (e.g. metal ions) and substrates are easily added to the reaction and the expression environment is highly controlled. This enables the production of fully active complex biocatalysts. The *E. coli*-based platform has successfully been used to produce enzymes with metal cofactors such as [FeFe] hydrogenases [36–40] and P450 monooxygenase [41]. Using multicopper oxidases as a

**Correspondence:** Prof. Michael C. Jewett, Department of Chemical and Biological Engineering, Northwestern University, Evanston, IL 60208, USA  
**E-mail:** m-jewett@northwestern.edu

**Abbreviations:** CFPS, Cell-free protein synthesis; MCO, Multicopper oxidase

model, we investigated the use of cell-free biosynthesis to produce copper enzymes.

Multicopper oxidases (MCOs) are a well-characterized group of enzymes that couple the reduction of dioxygen to water with the oxidation of a broad range of substrates [42]. Because of their broad activity, they have attracted significant attention for use in a variety of biotechnological applications including wastewater decolorization, pulp delignification, and enzymatic fuel cells [43]. Natively, MCOs are found in all kingdoms of life and are known to function in plant development, metal homeostasis, and fungal delignification [44]. To perform their function, MCOs require four copper atoms arranged in two distinct metal centers, a mononuclear type 1 copper and a trinuclear copper cluster formed by a type 2 copper and a pair of type 3 coppers [45]. Substrates are oxidized at the type 1 copper center and electrons are then transferred to the trinuclear copper cluster where oxygen is reduced to water [45].

MCOs were chosen for this study because their reported in vivo expression yields tend to be low and copper incorporation can be challenging [46]. In the present work, two bacterial MCOs from *Nitrosomonas europaea* (NeMCO) and *Streptomyces coelicolor* (ScMCO) and two archaeal MCOs from *Haloarcula marismortui* (HmMCO) and *Nitrosopumilus maritimus* (NmMCO) were synthesized. Using the *E. coli* CFPS platform, three of the four MCOs had final yields of more than 1 mg mL<sup>-1</sup>. Significant amounts of copper (100 μM) can be present during synthesis without impacting expression levels; however, given the high expression levels of the MCO this is not sufficient to completely metal load the protein. We found that CuSO<sub>4</sub> stimulates MCO activity by post-translational addition after CFPS. To the best of our knowledge, this is the first report on cell-free synthesis of an active MCO in the *E. coli*-based system. Our results, therefore, expand the ever-growing utility of CFPS techniques to include the production of MCOs and proteins with copper cofactors.

## 2 Materials and methods

### 2.1 Bacterial strains and plasmids

Highly active S30 crude extracts were generated from a genomically recoded release factor 1 (RF1) deficient *E. coli* strain (*E. coli* C321.ΔA.705) [47]. Multicopper oxidase genes from *Haloarcula marismortui* (HmMCO, accession no. WP\_004594571), *Nitrosopumilus maritimus* (NmMCO, accession no. WP\_012215737) and *Streptomyces coelicolor* (ScMCO, accession no. WP\_003972284) were codon optimized with *E. coli* class 2 codons, native secretion signals were omitted, and synthesized as gBlocks® (IDT Inc., Coralville, IA, USA). The *Nitrosomonas europaea* gene (NeMCO accession no. WP\_011111538)

was not codon optimized, but it was also synthesized using gBlocks®. The gBlocks® were incorporated into the vector pY71 using Gibson Assembly [48] and the DNA sequence verified (Eurofins Genomics, Huntsville, AL, USA). The four MCO genes were cloned in frame with a C-terminal tag comprised sequentially of a TEV cleavage site, (Gly<sub>3</sub>Ser)<sub>2</sub> linker, β-strand 11 of GFP [49], a Gly<sub>3</sub>Ser linker and a Strep-tag. The GFP11 sequence was inserted before the Strep-tag to facilitate a future study for high-throughput screening soluble expression of MCO variants. The following plasmids were tested in in vitro CFPS assays: pY71-NeMCO-TGS, pY71-HmMCO-TGS, pY71-NmMCO-TGS and pY71-ScMCO-TGS.

### 2.2 Preparation of cell extracts

*E. coli* cells were grown in 1 L of 2xYTPG (yeast extract 10 g L<sup>-1</sup>, tryptone 16 g L<sup>-1</sup>, NaCl 5 g L<sup>-1</sup>, K<sub>2</sub>HPO<sub>4</sub> 7 g L<sup>-1</sup>, KH<sub>2</sub>PO<sub>4</sub> 3 g L<sup>-1</sup> and glucose 18 g L<sup>-1</sup>, pH 7.2) in a 2.5-L Tunair flask (IBI Scientific, Peosta, IA) at 34°C and 220 rpm with inoculation of 20 mL overnight cultures (initial OD<sub>600</sub> of ~0.05). Cell growth, collection and extracts were prepared as described by Kwon and Jewett [50].

### 2.3 CFPS reactions

Standard CFPS reactions were performed using the PANOX-SP system as previously described [51]. For copper loading, CuSO<sub>4</sub> was added to the standard system with final concentrations of 10, 20, 50, 100, 200 and 500 μM. For scale-up reactions with 50 and 100 μL mixture, the reactions were carried out in 2 mL microcentrifuge tubes. All CFPS reactions were incubated for 20 h at 30°C.

### 2.4 Quantification of synthesized MCOs

The yields of synthesized MCOs were quantified by the incorporation of <sup>14</sup>C-leucine into trichloroacetic acid-precipitable radioactivity using a liquid scintillation counter (MicroBeta2, PerkinElmer, Waltham, MA) as described previously [52]. Total protein yields were measured directly after cell-free reactions with the 15 μL mixture. To determine the soluble protein yield, 15 μL of the reaction mixture was centrifuged at 12 000 ×g and 4°C for 15 min [53]. 10 μL of the supernatant were taken out for soluble MCO quantification.

### 2.5 Autoradiography analysis

After the cell-free reaction, 3 μL of each sample was loaded on a 4–12% NuPAGE SDS-PAGE gel (Invitrogen). The gel was stained using SimplyBlue™ SafeStain solution (Invitrogen) and destained in water. Then, the gel was fixed with cellophane films (Bio-Rad), dried overnight in a GelAir Dryer (Bio-Rad) without heating, and exposed for 48 h on a Storage Phosphor Screen (GE Healthcare)

Biosciences, Pittsburgh, PA). The autoradiogram was scanned using a Storm Imager (GE Healthcare Biosciences, Pittsburgh, PA) and analyzed using Quantity One software (Bio-Rad, Hercules, CA).

## 2.6 MCO activity assay

MCO activity was measured colorimetrically by following oxidation of ABTS ( $\epsilon = 36.8 \text{ mM}^{-1}\text{cm}^{-1}$  at 405 nm), which turns blue upon oxidation [54]. Colorimetric assays were performed in 1 mL of 100 mM Britton-Robinson buffer at pH 4.0 containing 4 mM ABTS using an Agilent 4853 spectrophotometer. All cell-free reactions were diluted 10-fold and incubated with 0 to 5 mM  $\text{CuSO}_4$  or 0 to 500  $\mu\text{M}$   $\text{Cu}(\text{CH}_3\text{CN})_4\text{PF}_6$  and at room temperature for 2 h. In cases where  $\text{Cu}(\text{CH}_3\text{CN})_4\text{PF}_6$  was added, the reaction was transferred into an anaerobic Coy chamber and allowed to equilibrate for 2 h prior to addition of copper. All reactions were initiated by addition of 40 to 80  $\mu\text{L}$  of diluted cell-free samples and reported values are averaged from at least three replicates. All reactions were performed at room temperature.

## 2.7 In vivo expression of MCOs


MCOs were expressed in *E. coli* strain BL21(DE3) and grown in homemade autoinduction media [55]. Cells were grown at 37°C to an  $\text{OD}_{600}$  of ~0.6 and then moved to 20°C for 16 h. Prior to harvest cells were incubated with 5 mM  $\text{CuSO}_4$  for 2 h and then centrifuged for 10 min at 6 400  $\times g$  and 4°C. Cells were suspended in 50 mM TRIS pH 8.0 and 500 mM NaCl and lysed by sonication for 8 min with 1 s

pulses and 3 s resting time. Lysates were centrifuged for 1 h at 185 000  $\times g$  and 4°C and the supernatants applied to a streptactin column (IBA Life Sciences). MCOs were eluted with 50 mM TRIS pH 8.0, 500 mM NaCl, and 5 mM *d*-desthiobiotin and then concentrated in Amicon centrifugal devices with a 30 kDa nominal molecular weight cutoff (EMD Millipore). The concentrated MCOs were then further purified on a Superdex 200 Increase (GE Healthcare Life Sciences) size exclusion column equilibrated with 50 mM TRIS pH 8.0 and 500 mM NaCl. Final yields of the MCOs were determined using their extinction coefficients at 280 nm.

## 3 Results and discussion

### 3.1 CFPS enables soluble high-level expression of four MCOs

The *E. coli*-based cell-free system used in this work is similar to a previously published report [51], with only minor modifications (see above). The reactions were allowed to run for 20 h in batch operation and the yields of cell-free synthesized MCOs were quantified by monitoring  $^{14}\text{C}$ -leucine incorporation. As shown in Fig. 1A, the best expressing variant, NeMCO, gave a protein yield of  $1.2 \pm 0.1 \text{ mg mL}^{-1}$  (>95% soluble), while ScMCO yields reached less than 0.2 mg  $\text{mL}^{-1}$ . HmMCO and NmMCO also reached high final yields of ~1 mg  $\text{mL}^{-1}$ ; however, their soluble fractions were lower than NeMCO, constituting 36 and 70% of the total protein expression, respectively. These yields are higher than those typically



**Figure 1.** Cell-free protein synthesis of two bacterial and two archaeal MCOs. (A) Protein yields of MCOs. T, total protein; S, soluble protein. Error bars represent standard deviations from three independent samples. (B) SDS-PAGE analysis of cell-free produced MCOs. Each MCO is indicated by an asterisk. Expected molecular weight: NeMCO (41.1 kDa), HmMCO (43.5 kDa), NmMCO (51.6 kDa), ScMCO (39.4 kDa). M, protein molecular weight marker; NC, negative control without plasmid in the reaction. (C) Autoradiogram of radioactive labeled proteins.

**Table 1.** Examples of MCOs heterologously expressed in vivo

MCO <sup>a)</sup>	Source	Expression host	Titer (mg L <sup>-1</sup> )	Reference
NeMCO	<i>Nitrosomonas europaea</i>	<i>E. coli</i>	42.0 ± 14 <sup>b)</sup>	This Study
HmMCO	<i>Haloarcula marismortui</i>	<i>E. coli</i>	Inclusion bodies <sup>b)</sup>	This Study
NmMCO	<i>Nitrosopumilus maritimus</i>	<i>E. coli</i>	Inclusion bodies <sup>b)</sup>	This Study
ScMCO	<i>Streptomyces coelicolor</i>	<i>E. coli</i>	11.0 ± 4.1 <sup>b)</sup>	This Study
Laccase	<i>Bacillus subtilis</i>	<i>E. coli</i>	20	[56]
Laccase	<i>Aeromonas hydrophila</i>	<i>E. coli</i>	10	[57]
Laccase	<i>Streptomyces coelicolor</i>	<i>Streptomyces lividans</i>	350	[63]
Laccase	<i>Myceliophthora thermophila</i>	<i>Saccharomyces cerevisiae</i>	18	[64]
Laccase	<i>Schizophyllum commune</i>	<i>Pycnoporus cinnabarinus</i>	1200	[65]

a) Laccase belongs to the multicopper oxidase family (MCO).

b) Yield after metal loading and growth in autoinduction media. In all cases significant quantities of inclusion bodies were observed in addition to the reported soluble expression yields.

observed in vivo. For example, the expression levels of laccases (belonging to the MCO family) from *Bacillus subtilis* and *Aeromonas hydrophila* in *E. coli* are ~20 mg L<sup>-1</sup> and ~10 mg L<sup>-1</sup>, respectively [56, 57]. The fact that different MCOs from different organisms expressed at different titers was not surprising, as expression levels are known to be highly dependent on several gene sequence features, such as the position dependent codon usage, ribosome binding site, mRNA structure, etc. [58]. Even with different titers, expressed MCOs could be identified from the *E. coli* endogenous proteins using SDS-PAGE gels. These results show that the correct molecular weights were obtained (Fig. 1B) and were further confirmed by autoradiogram analysis (Fig. 1C). Because NeMCO produced the highest soluble yields, subsequent analysis focused on it as a target.


To directly compare our CFPS expression system to conventional in vivo strategies, the four MCO constructs were also expressed in living *E. coli* (Table 1). In vivo expression yields for all constructs were less than 50 mg L<sup>-1</sup>, which is lower than that obtained with the *E. coli*-based CFPS system. Notably, two of the constructs, HmMCO and NmMCO, form only inclusion bodies in vivo, whereas they are soluble in vitro. This highlights a potential advantage for CFPS. We believe the increased solubility in vitro may arise from the dilute total protein environment in the extract (~10 mg mL<sup>-1</sup> total *E. coli* protein) as compared to the cytoplasm (~200–300 mg mL<sup>-1</sup> total *E. coli* protein), which helps favor protein folding over aggregation.

### 3.2 Effect of scalability and copper concentration on CFPS yields of NeMCO

To demonstrate the potential application of CFPS of MCOs, the reaction volume of NeMCO was scaled from 15 to 100 µL. While the reaction volume increased more than six times compared to the standard reaction, NeMCO expression levels were similar to small scale reactions with total yields of ~1.2 mg mL<sup>-1</sup> (Fig. 2A). Importantly,

approximately 95% of the expressed protein remained soluble.

Copper is necessary for reconstitution of the MCO's activity. To this end, the effect of copper concentration



**Figure 2.** (A) Cell-free protein synthesis of NeMCO with scale-up systems. T, total NeMCO; S, soluble NeMCO. (B) Effect of Cu<sup>2+</sup> on NeMCO synthesis in the cell-free system. Error bars represent standard deviations from three independent samples.




on expression levels was studied using NeMCO as copper is toxic for many bacteria. At moderate concentrations, 0 to 100  $\mu\text{M}$ , there is no obvious decrease in protein yield (Fig. 2B), suggesting some benefits for CFPS avoiding toxicity that occurs in cells. However, at high concentrations of copper, 200  $\mu\text{M}$  and above, expression drops significantly and copper appears to be toxic to the combined transcription-translation system. Consistent with the hypothesis that copper may be toxic to CFPS, we found that copper concentrations of 100  $\mu\text{M}$  reduced expression of superfolder green fluorescent protein (which does not have copper cofactors) four-fold as compared to reactions without supplemented copper (data not shown). Our results suggest that copper appears to be toxic to CFPS at high concentrations, but less toxic when manufacturing copper-containing proteins, which likely sequester the copper.

### 3.3 Cell-free synthesized NeMCO is active, and stimulated by post-translational copper addition

To verify that the cell-free synthesized NeMCO was active, assays were performed by following the oxidation of 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS), a non-native substrate, which is blue in its oxidized form. Initially these experiments focused on NeMCO that was synthesized in the presence of 100  $\mu\text{M}$  copper. Our results indicate that synthesis in 100  $\mu\text{M}$  copper is not sufficient to obtain active protein but post-translational addition of  $\text{CuSO}_4$  stimulates activity (Fig. 3). In cell-free reactions the final concentration of NeMCO is  $\sim 30 \mu\text{M}$  and each monomer requires at least four coppers. It is therefore not unexpected that NeMCO is not active immediately following the cell-free reaction since there is not enough copper in solution to fully load the enzyme. To test if post-translational addition of copper stimulates activity, reactions were diluted and Cu(II) in the form of  $\text{CuSO}_4$  was added (Fig. 3). Our data show that activity is stimulated by relatively low concentrations of copper and is diminished if too much copper is added. Diminished activity is likely due to protein precipitation caused by excess copper ions. Importantly, we do not observe copper acting as a catalyst alone to oxidize the substrate. Negative controls with similarly prepared CFPS reactions without plasmids were performed with and without copper and oxidation of the dye was not observed.

Previous reports suggest that maximal activity of MCOs is obtained when copper is loaded as Cu(I) [56]. To test this possibility, cell-free reactions were taken into an anaerobic Coy chamber and incubated with a Cu(I) compound,  $\text{Cu}(\text{CH}_3\text{CN})_4\text{PF}_6$ , in a similar fashion as the  $\text{CuSO}_4$  experiments. These data show that incubating with Cu(I) versus Cu(II) results in similar levels of activity. Cu(I) could not be tested at concentrations above 500  $\mu\text{M}$  because at these concentrations direct reduction of ABTS



**Figure 3.** Effect of copper concentration on specific activity. White, no additional copper added; light gray, addition of Cu(I) as  $\text{Cu}(\text{CH}_3\text{CN})_4\text{PF}_6$ ; dark gray, addition of Cu(II) as  $\text{CuSO}_4$ . Error bars represent standard deviations from three independent samples.

by unbound Cu(I) is observed preventing reliable data collection. Unfortunately, the current specific activity of NeMCO is  $\sim 10\%$  the rate of the same protein produced in vivo lacking a C-terminal tag [59]. We attribute this diminished rate to the large C-terminally encoded tag, which has been previously shown to greatly reduce MCO activity [60]. Indeed, the lower specific activity in the CFPS system matches the activity of in vivo expressed NeMCO in *E. coli* with the same C-terminal tag (data not shown). Thus, our work provides a promising route to discovery-based applications that require rapid, low-cost protein expression, such as high-throughput screening of soluble MCO variants.

## 4 Concluding remarks: CFPS offers a new approach for producing MCOs at high titers

In conclusion, multiple MCOs were expressed in an *E. coli*-based CFPS system with high yields ( $>1 \text{ mg mL}^{-1}$ ). Scale-up reactions were achieved from 15 to 100  $\mu\text{L}$  without decreasing productivity and solubility. Importantly, MCO activity can be obtained by simple addition of copper ions post-translation and our work also establishes a threshold for copper tolerance in *E. coli*-based CFPS. Our work is important because it allows for expression levels that are adequate for testing MCO activity, setting the stage for a high-throughput expression platform that can be executed in a day through the use of PCR templates, as compared to days to weeks in vivo [6].

Indeed, our approach may provide some potential advantages for high yielding expression in high-throughput as compared to existing state of the art. As summarized in the review paper [61], different heterologous

hosts have been used to express MCOs (mostly laccases) including bacteria systems (e.g. *E. coli* and *Streptomyces lividans*), yeast systems (e.g. *Saccharomyces cerevisiae*, *Pichia pastoris*, etc.), and filamentous fungi systems (e.g. *Aspergillus niger*, *Trichoderma reesei*, etc.). Previous expression titers are typically lower than those observed here (~10–50 mg L<sup>-1</sup>). A summary of these results, including the in vivo expression of the constructs expressed in vitro in this study, can be found in Table 1.

Looking forward, opportunities to improve the CFPS system include gaining a better understanding of the copper loading conditions, enabling efficient glycosylation that is required for some archaeal enzymes [62], and demonstrating higher preparative reaction scales of active MCOs. However, we note that the *E. coli* CFPS system has already been shown to scale linearly from µL reactions to the 100 L scale, an expansion factor of 10<sup>6</sup>, to enable manufacturing scale synthesis of complex proteins [26]. We therefore anticipate that CFPS systems will provide a new avenue for producing MCOs in the future.


*This work was supported by the Advanced Research Projects Agency-Energy (ARPA-E) REMOTE program (DE-AR0000435).*

*The authors declare no financial or commercial conflict of interest.*

## 5 References

- Hodgman, C. E., Jewett, M. C., Cell-free synthetic biology: Thinking outside the cell. *Metab. Eng.* 2012, 14, 261–269.
- Swartz, J. R., Transforming biochemical engineering with cell-free biology. *AIChE J.* 2012, 58, 513.
- Albayrak, C., Swartz, J. R., Cell-free co-production of an orthogonal transfer RNA activates efficient site-specific non-natural amino acid incorporation. *Nucleic Acids Res.* 2013, 41, 5949–5963.
- Anderson, M. J., Stark, J. C., Hodgman, C. E., Jewett, M. C., Energizing eukaryotic cell-free protein synthesis with glucose metabolism. *FEBS Lett.* 2015, 589, 1723–1727.
- Bundy, B. C., Swartz, J. R., Site-specific incorporation of *p*-propargyloxyphenylalanine in a cell-free environment for direct protein–protein click conjugation. *Bioconjug. Chem.* 2010, 21, 255–263.
- Carlson, E. D., Gan, R., Hodgman, C. E., Jewett, M. C., Cell-free protein synthesis: Applications come of age. *Biotechnol. Adv.* 2012, 30, 1185–1194.
- Caschera, F., Noireaux, V., Synthesis of 2.3 mg/ml of protein with an all *Escherichia coli* cell-free transcription–translation system. *Biochimie* 2014, 99, 162–168.
- Choudhury, A., Hodgman, C. E., Anderson, M. J., Jewett, M. C., Evaluating fermentation effects on cell growth and crude extract metabolic activity for improved yeast cell-free protein synthesis. *Biochem. Eng. J.* 2014, 91, 140–148.
- Gan, R., Jewett, M. C., A combined cell-free transcription–translation system from *Saccharomyces cerevisiae* for rapid and robust protein synthesis. *Biotechnol. J.* 2014, 9, 641–651.
- Goshima, N., Kawamura, Y., Fukumoto, A., Miura, A. et al., Human protein factory for converting the transcriptome into an in vitro-expressed proteome. *Nat. Methods* 2008, 5, 1011–1017.
- Heinzelman, P., Schoborg, J. A., Jewett, M. C., pH responsive granulocyte colony-stimulating factor variants with implications for treating Alzheimer’s disease and other central nervous system disorders. *Protein Eng. Des. Sel.* 2015, DOI:10.1093/protein/gzv022.
- Hodgman, C. E., Jewett, M. C., Optimized extract preparation methods and reaction conditions for improved yeast cell-free protein synthesis. *Biotechnol. Bioeng.* 2013, 110, 2643–2654.
- Hodgman, C. E., Jewett, M. C., Characterizing IGR IRES-mediated translation initiation for use in yeast cell-free protein synthesis. *New Biotechnol.* 2014, 31, 499–505.
- Hong, S. H., Kwon, Y. C., Jewett, M. C., Non-standard amino acid incorporation into proteins using *Escherichia coli* cell-free protein synthesis. *Front. Chem.* 2014, 2, 34.
- Hong, S. H., Kwon, Y. C., Martin, R. W., Des Soye, B. J. et al., Improving cell-free protein synthesis through genome engineering of *Escherichia coli* lacking release factor 1. *ChemBioChem* 2015, 16, 844–853.
- Kanter, G., Yang, J., Voloshin, A., Levy, S. et al., Cell-free production of scFv fusion proteins: An efficient approach for personalized lymphoma vaccines. *Blood* 2007, 109, 3393–3399.
- Kwon, Y. C., Lee, K. H., Kim, H. C., Han, K. et al., Cloning-independent expression and analysis of ω-transaminases by use of a cell-free protein synthesis system. *Appl. Environ. Microbiol.* 2010, 76, 6295–6298.
- Quast, R. B., Claussnitzer, I., Merk, H., Kubick, S., Gerrits, M., Synthesis and site-directed fluorescence labeling of azido proteins using eukaryotic cell-free orthogonal translation systems. *Anal. Biochem.* 2014, 451, 49.
- Sachse, R., Dondapati, S. K., Fenz, S. F., Schmidt, T., Kubick, S., Membrane protein synthesis in cell-free systems: From bio-mimetic systems to bio-membranes. *FEBS Lett.* 2014, 588, 2774–2781.
- Schoborg, J. A., Hodgman, C. E., Anderson, M. J., Jewett, M. C., Substrate replenishment and byproduct removal improve yeast cell-free protein synthesis. *Biotechnol. J.* 2014, 9, 630–640.
- Stafford, R. L., Matsumoto, M. L., Yin, G., Cai, Q. et al., In vitro Fab display: A cell-free system for IgG discovery. *Protein Eng. Des. Sel.* 2014, 27, 97–109.
- Uhlemann, E. M. E., Pierson, H. E., Fillingame, R. H., Dmitriev, O. Y., Cell-free synthesis of membrane subunits of ATP synthase in phospholipid bicelles: NMR shows subunit a fold similar to the protein in the cell membrane. *Protein Sci.* 2012, 21, 279–288.
- Yang, J., Kanter, G., Voloshin, A., Levy, R., Swartz, J. R., Expression of active murine granulocyte-macrophage colony-stimulating factor in an *Escherichia coli* cell-free system. *Biotechnol. Prog.* 2004, 20, 1689–1696.
- Yin, G., Garces, E. D., Yang, J., Zhang, J. et al., Aglycosylated antibodies and antibody fragments produced in a scalable in vitro transcription–translation system. *MAbs* 2012, 4, 217–225.
- Yin, G., Swartz, J. R., Enhancing multiple disulfide bonded protein folding in a cell-free system. *Biotechnol. Bioeng.* 2004, 86, 188–195.
- Zawada, J. F., Yin, G., Steiner, A. R., Yang, J. et al., Microscale to manufacturing scale-up of cell-free cytokine production – a new approach for shortening protein production development timelines. *Biotechnol. Bioeng.* 2011, 108, 1570–1578.
- Kim, T. W., Chokhawala, H. A., Hess, M., Dana, C. M. et al., High-throughput in vitro glycoside hydrolase (HIGH) screening for enzyme discovery. *Angew. Chem. Int. Ed.* 2011, 50, 11215–11218.
- Makino, S., Beebe, E. T., Markley, J. L., Fox, B. G., Cell-free protein synthesis for functional and structural studies. *Methods Mol. Biol.* 2014, 1091, 161–178.

- [29] Bujara, M., Schümperli, M., Billerbeck, S., Heinemann, M., Panke, S., Exploiting cell-free systems: Implementation and debugging of a system of biotransformations. *Biotechnol. Bioeng.* 2010, *106*, 376–389.
- [30] Chappell, J., Jensen, K., Freemont, P. S., Validation of an entirely in vitro approach for rapid prototyping of DNA regulatory elements for synthetic biology. *Nucleic Acids Res.* 2013, *41*, 3471–3481.
- [31] Chappell, J., Takahashi, M. K., Lucks, J. B., Creating small transcription activating RNAs. *Nat. Chem. Biol.* 2015, *11*, 214–220.
- [32] Pardee, K., Green, A. A., Ferrante, T., Cameron, D. E. et al., Paper-based synthetic gene networks. *Cell* 2014, *159*, 940–954.
- [33] Shin, J., Noireaux, V., An *E. coli* cell-free expression toolbox: Application to synthetic gene circuits and artificial cells. *ACS Synth. Biol.* 2012, *1*, 29–41.
- [34] Siegal-Gaskins, D., Tuza, Z. A., Kim, J., Noireaux, V., Murray, R. M., Gene circuit performance characterization and resource usage in a cell-free “breadboard”. *ACS Synth. Biol.* 2014, *3*, 416–425.
- [35] Sun, Z. Z., Yeung, E., Hayes, C. A., Noireaux, V., Murray, R. M., Linear DNA for rapid prototyping of synthetic biological circuits in an *Escherichia coli* based TX-TL cell-free system. *ACS Synth. Biol.* 2014, *3*, 387–397.
- [36] Boyer, M. E., Stapleton, J. A., Kuchenreuther, J. M., Wang, C. W., Swartz, J. R., Cell-free synthesis and maturation of [FeFe]hydrogenases. *Biotechnol. Bioeng.* 2008, *99*, 59–67.
- [37] Kuchenreuther, J. M., Britt, R. D., Swartz, J. R., New insights into [FeFe]hydrogenase activation and maturase function. *PLoS One* 2012, *7*, e45850.
- [38] Kuchenreuther, J. M., George, S. J., Grady-Smith, C. S., Cramer, S. P., Swartz, J. R., Cell-free H-cluster synthesis and [FeFe] hydrogenase activation: All five CO and CN<sup>-</sup> ligands derive from tyrosine. *PLoS One* 2011, *6*, e20346.
- [39] Kuchenreuther, J. M., Myers, W. K., Stich, T. A., George, S. J. et al., A radical intermediate in tyrosine scission to the CO and CN<sup>-</sup> ligands of FeFe hydrogenase. *Science* 2013, *342*, 472–475.
- [40] Kuchenreuther, J. M., Myers, W. K., Suess, D. L. M., Stich, T. A. et al., The HydG enzyme generates an Fe(CO)<sub>2</sub>(CN) synthon in assembly of the FeFe hydrogenase H-cluster. *Science* 2014, *343*, 424–427.
- [41] Kwon, Y. C., Oh, I. S., Lee, N., Lee, K. H. et al., Integrating cell-free biosyntheses of heme prosthetic group and apoenzyme for the synthesis of functional P450 monooxygenase. *Biotechnol. Bioeng.* 2013, *110*, 1193–1200.
- [42] Nakamura, K., Go, N., Function and molecular evolution of multicopper blue proteins. *Cell. Mol. Life Sci.* 2005, *62*, 2050–2066.
- [43] Kunamneni, A., Plou, F. J., Ballesteros, A., Alcalde, M., Laccases and their applications: A patent review. *Recent Pat. Biotechnol.* 2008, *2*, 10–24.
- [44] Mayer, A. M., Staples, R. C., Laccase: New functions for an old enzyme. *Phytochemistry* 2002, *60*, 551–565.
- [45] Solomon, E. I., Sundaram, U. M., Machonkin, T. E., Multicopper oxidases and oxygenases. *Chem. Rev.* 1996, *96*, 2563–2606.
- [46] Santhanam, N., Vivanco, J. M., Decker, S. R., Reardon, K. F., Expression of industrially relevant laccases: Prokaryotic style. *Trends Biotechnol.* 2011, *29*, 480–489.
- [47] Lajoie, M. J., Rovner, A. J., Goodman, D. B., Aerni, H. R. et al., Genomically recoded organisms expand biological functions. *Science* 2013, *342*, 357–360.
- [48] Gibson, D. G., Young, L., Chuang, R. Y., Venter, J. C. et al., Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat. Methods* 2009, *6*, 343–345.
- [49] Cabantous, S., Waldo, G. S., In vivo and in vitro protein solubility assays using split GFP. *Nat. Methods* 2006, *3*, 845–854.
- [50] Kwon, Y. C., Jewett, M. C., High-throughput preparation methods of crude extract for robust cell-free protein synthesis. *Sci. Rep.* 2015, *5*, 8663.
- [51] Jewett, M. C., Swartz, J. R., Mimicking the *Escherichia coli* cytoplasmic environment activates long-lived and efficient cell-free protein synthesis. *Biotechnol. Bioeng.* 2004, *86*, 19–26.
- [52] Jewett, M. C., Swartz, J. R., Rapid expression and purification of 100 nmol quantities of active protein using cell-free protein synthesis. *Biotechnol. Prog.* 2004, *20*, 102–109.
- [53] Jewett, M. C., Calhoun, K. A., Voloshin, A., Wu, J. J., Swartz, J. R., An integrated cell-free metabolic platform for protein production and synthetic biology. *Mol. Syst. Biol.* 2008, *4*, 220.
- [54] Lawton, T. J., Rosenzweig, A. C., Detection and characterization of a multicopper oxidase from *Nitrosomonas europaea*. *Methods Enzymol.* 2011, *496*, 423–433.
- [55] Studier, F. W., Protein production by auto-induction in high-density shaking cultures. *Protein Expr. Purif.* 2005, *41*, 207–234.
- [56] Durão, P., Chen, Z., Fernandes, A. T., Hildebrandt, P. et al., Copper incorporation into recombinant CotA laccase from *Bacillus subtilis*: Characterization of fully copper loaded enzymes. *J. Biol. Inorg. Chem.* 2008, *13*, 183–193.
- [57] Ng, I. S., Zhang, X., Zhang, Y., Lu, Y., Molecular cloning and heterologous expression of laccase from *Aeromonas hydrophila* NIU01 in *Escherichia coli* with parameters optimization in production. *Appl. Biochem. Biotechnol.* 2013, *169*, 2223–2235.
- [58] Hockenberry, A. J., Sirer, M. I., Amaral, L. A. N., Jewett, M. C., Quantifying position-dependent codon usage bias. *Mol. Biol. Evol.* 2014, *31*, 1880–1893.
- [59] Lawton, T. J., Sayavedra-Soto, L. A., Arp, D. J., Rosenzweig, A. C., Crystal structure of a two-domain multicopper oxidase: Implications for the evolution of multicopper proteins. *J. Biol. Chem.* 2009, *284*, 10174–10180.
- [60] Autore, F., Del Vecchio, C., Fraternali, F., Giardina, P. et al., Molecular determinants of peculiar properties of a *Pleurotus ostreatus* laccase: Analysis by site-directed mutagenesis. *Enzyme Microb. Technol.* 2009, *45*, 507–513.
- [61] Piscitelli, A., Pezzella, C., Giardina, P., Faraco, V., Sannia, G., Heterologous laccase production and its role in industrial applications. *Bioeng. Bugs* 2010, *1*, 254–264.
- [62] Guarino, C., DeLisa, M. P., A prokaryote-based cell-free translation system that efficiently synthesizes glycoproteins. *Glycobiology* 2012, *22*, 596–601.
- [63] Dubé, E., Shareck, F., Hurtubise, Y., Daneault, C., Beaugard, M., Homologous cloning, expression, and characterisation of a laccase from *Streptomyces coelicolor* and enzymatic decolourisation of an indigo dye. *Appl. Microbiol. Biotechnol.* 2008, *79*, 597–603.
- [64] Bulter, T., Alcalde, M., Sieber, V., Meinhold, P. et al., Functional expression of a fungal laccase in *Saccharomyces cerevisiae* by directed evolution. *Appl. Environ. Microbiol.* 2003, *69*, 987–995.
- [65] Alves, A. M. C. R., Record, E., Lomascolo, A., Scholtmeijer, K. et al., Highly efficient production of laccase by the basidiomycete *Pycnoporus cinnabarinus*. *Appl. Environ. Microbiol.* 2004, *70*, 6379–6384.



## Biotechnology Journal – list of articles published in the February 2016 issue.

### Editorial

#### Transforming biotechnology with synthetic biology

George Guo-Qiang Chen and Michael C. Jewett

<http://dx.doi.org/10.1002/biot.201600010>

### BTJ-Forum

#### In Memoriam of Prof. Bernard Witholt

Manfred Zinn, Sang Yup Lee and George Guo-Qiang Chen

<http://dx.doi.org/10.1002/biot.201500096>

### BTJ-Forum

#### Meeting Report:

#### Cold Spring Harbor Asia Synthetic Biology Meeting

Ivan Hajnal

<http://dx.doi.org/10.1002/biot.201400836>

### Review

#### Minimal genome:

#### Worthwhile or worthless efforts toward being smaller?

Donghui Choe, Suhyung Cho, Sun Chang Kim and Byung-Kwan Cho

<http://dx.doi.org/10.1002/biot.201400838>

### Research Article

#### Cell-free protein synthesis enables high yielding synthesis of an active multicopper oxidase

Jian Li, Thomas J. Lawton, Jan S. Kostecki, Alex Nisthal, Jia Fang, Stephen L. Mayo, Amy C. Rosenzweig and Michael C. Jewett

<http://dx.doi.org/10.1002/biot.201500030>

### Research Article

#### Engineering of core promoter regions enables the construction of constitutive and inducible promoters in *Halomonas* sp.

Tingting Li, Teng Li, Weiyue Ji, Qiuyue Wang, Haoqian Zhang, Guo-Qiang Chen, Chunbo Lou and Qi Ouyang

<http://dx.doi.org/10.1002/biot.201400828>

### Research Article

#### Genome mining of astaxanthin biosynthetic genes from *Sphingomonas* sp. ATCC 55669 for heterologous overproduction in *Escherichia coli*

Tian Ma, Yuanjie Zhou, Xiaowei Li, Fayin Zhu, Yongbo Cheng, Yi Liu, Zixin Deng and Tiangang Liu

<http://dx.doi.org/10.1002/biot.201400827>

### Research Article

#### A cell-free expression and purification process for rapid production of protein biologics

Challise J. Sullivan, Erik D. Pendleton, Henri H. Sasmor, William L. Hicks, John B. Farnum, Machiko Muto, Eric M. Amendt, Jennifer A. Schoborg, Rey W. Martin, Lauren G. Clark, Mark J. Anderson, Alaksh Choudhury, Raffaella Fior, Yu-Hwa Lo, Richard H. Griffey, Stephen A. Chappell, Michael C. Jewett, Vincent P. Mauro and John Dresios

<http://dx.doi.org/10.1002/biot.201500214>

### Research Article

#### Co-production of hydrogen and ethanol from glucose by modification of glycolytic pathways in *Escherichia coli* – from Embden-Meyerhof-Parnas pathway to pentose phosphate pathway

Eunhee Seol, Balaji Sundara Sekar, Subramanian Mohan Raj and Sunghoon Park

<http://dx.doi.org/10.1002/biot.201400829>

### Research Article

#### A fluorescein-labeled AmpC $\beta$ -lactamase allows rapid characterization of $\beta$ -lactamase inhibitors by real-time fluorescence monitoring of the $\beta$ -lactamase-inhibitor interactions

Man-Wah Tsang, Pak-Ho Chan, Sze-Yan Liu, Kwok-Yin Wong and Yun-Chung Leung

<http://dx.doi.org/10.1002/biot.201400861>



Research Article

**Alphavirus capsid proteins self-assemble into core-like particles in insect cells: A promising platform for nanoparticle vaccine development**

*Mia C. Hikke, Corinne Geertsema, Vincen Wu, Stefan W. Metz, Jan W. van Lent, Just M. Vlak and Gorben P. Pijlman*

<http://dx.doi.org/10.1002/biot.201500147>

Research Article

**Cell-free protein synthesis of a cytotoxic cancer therapeutic: Onconase production and a just-add-water cell-free system**

*Amin S. M. Salehi, Mark Thomas Smith, Anthony M. Bennett, Jacob B. Williams, William G. Pitt and Bradley C. Bundy*

<http://dx.doi.org/10.1002/biot.201500237>

Research Article

**Non-monotonic course of protein solubility in aqueous polymer-salt solutions can be modeled using the sol-mxDLVO model**

*Marcel Herhut, Christoph Brandenbusch and Gabriele Sadowski*

<http://dx.doi.org/10.1002/biot.201500123>

Biotech Method

**Rational plasmid design and bioprocess optimization to enhance recombinant adeno-associated virus (AAV) productivity in mammalian cells**

*Verena V. Emmerling, Antje Pegel, Ernest G. Milian, Alina Venereo-Sanchez, Marion Kunz, Jessica Wegele, Amine A. Kamen, Stefan Kochanek and Markus Hoerer*

<http://dx.doi.org/10.1002/biot.201500176>